

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/95152/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Ugakar, A.G., Kumar, B.C., Malapur, M.A. and Kerr, Andrew ORCID: <https://orcid.org/0000-0001-5569-4730> 2017. Petrography and geochemistry of Archaean Greywackes from Northern Part of the Dharwar-Shimoga Greenstone Belt, Western Dharwar Craton: implications for nature of provenance. Journal of the Geological Society of India 89 (5) , pp. 547-553. 10.1007/s12594-017-0643-6 file

Publishers page: <http://dx.doi.org/10.1007/s12594-017-0643-6>
<<http://dx.doi.org/10.1007/s12594-017-0643-6>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Petrography and Geochemistry of Archaean Greywackes from Northern Part of the Dharwar-Shimoga Greenstone Belt, Western Dharwar Craton: Implications for Nature of Provenance

A. G. UGARKAR

Department of Geology, Karnatak University, Dharwad 580 003, India.

CHANDAN KUMAR B. (Corresponding Author)

Department of Geology, Karnatak University, Dharwad 580 003, India.

*E-mail: rcgeo85@gmail.com

M. A. MALAPUR

Department of Geology, Karnatak University, Dharwad 580 003, India.

T. B. MANUVACHARI

Department of Geology, Karnatak University, Dharwad 580 003, India.

A.C. KERR

School of Earth and Ocean Sciences, Cardiff University, Cardiff, UK.

Acknowledgements: The authors are thankful to Drs. V. Balaram, D.S. Sarma and M. Ram Mohan of National Geophysical Research Institute, Hyderabad and Dr. G.R. Ravindra Kumar, Center for Earth Science Studies, Trivandrum for extending analytical facilities. This paper forms a part of study carried out under the Department of Science and Technology funded Major Research Project (ESS/16/343/2008) awarded to AGU. BCK acknowledges the financial support received from Council of Scientific Research-University Grants Commission in the form of Senior Research Fellowship (F.17-21/08(SA-1)). The authors thank the Chairman, Department of Studies in Geology, Karnatak University, Dharwad for providing infrastructure facilities to carry out the present study. The manuscript has been greatly improved by meticulous review, valuable comments and suggestions by an anonymous reviewer of the journal for which the authors are grateful.

Petrography and Geochemistry of Archaean Greywackes from Northern Part of the Dharwar-Shimoga Greenstone Belt, Western Dharwar Craton: Implications for Nature of Provenance

Abstract: Greywackes (Dharwar greywackes) are the most abundant rock types in the northern part of the Dharwar-Shimoga greenstone belt of the Western Dharwar Craton. They are distinctly immature rocks with poorly-sorted angular to sub-angular grains, comprising largely quartz, plagioclase feldspar and lithic fragments of volcanics (mafic+felsic), chert and quartzite, with subordinate biotite, K-feldspar and muscovite. They are characterized by almost uniform silica (59.78-67.96 wt%; av. 62.58), alkali (4.62-7.35 wt%; av. 5.41) contents, $\text{SiO}_2/\text{Al}_2\text{O}_3$ (3.71-5.25) ratios, and compositionally are comparable to the andesite and dacite. As compared to Ranibennur greywackes, located about 100 km south of Dharwad in the Dharwar-Shimoga greenstone belt, the Dharwar greywackes have higher K_2O , CaO , Zr , Y , ΣREE , Th/Sc , Zr/Cr , La/Sc and lower Sr , Cr , Ni , Sc , Cr/Th values. The chondrite normalized patterns of Dharwar greywackes are characterized by moderately fractionated REE patterns with moderate to high LREE enrichment, with almost flat HREE patterns and small negative Eu anomalies, suggesting felsic dominated source rocks in the provenance. The framework grains comprising felsic and mafic volcanics, feldspars and quartz suggest a mixed source in the provenance. The moderate CIA values ranging between 57 and 73, indicate derivation of detritus from fresh basement rocks and from nearby volcanic sources. The mixing calculations suggest that the average REE pattern is closely matching with a provenance having 40% dacite, 30% granite, 20% basalt and 10% tonalite. These greywackes were deposited in a subduction related forearc basin than a continental margin basin. Their La/Sc ratios are high (av. 4.07) compared to the Ranibennur greywackes (1.79), suggesting that the greywackes of the northern part of the basin received detritus from a more evolved continental crust than the greywackes of the central part of the Dharwar-Shimoga basin.

Keywords: Dharwar-Shimoga greenstone belt, Greywackes, Western Dharwar Craton,

Provenance, Geodynamic setting

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

Among clastic sediments, greywackes constitute the most abundant rocks of the Archaean sedimentation. As they retain significant concentrations of a variety of trace elements due to their diverse framework mineralogy and matrix characteristics, they are the extensively studied clastic metasediments of the greenstone belts all over the world (Taylor and McLennan, 1985; Naqvi et al., 1988; Srinivasan et al., 1989; Feng and Kerrich, 1990; Manikyamba et al., 1997; Ugarkar and Nyamati, 2002; Ootes et al., 2009; Devaraju et al., 2010; Absar and Sreenivas, 2015). The overall abundance of any clastic sedimentary component in the greenstone belts is of considerable significance. For example, the occurrence of detrital sediments like greywackes has a clear significance in terms of the then prevailing exogenic processes and environment. Such sediments contain important information about the source compositions, weathering conditions, nature of provenance and geodynamic settings of the depositional basins. In the Neoarchaean Dharwar-Shimoga greenstone belt of the Western Dharwar Craton greywackes are the most abundant clastic metasediments. In this paper, we present and discuss the petrography and geochemistry of greywackes from the northern part (Dharwar greywackes) of the Dharwar-Shimoga greenstone belt, with an aim of understanding sedimentation environment and nature of the provenance in the northern part of the Western Dharwar Craton.

GEOLOGY

The Dharwar-Shimoga greenstone belt of the well-known Western Dharwar Craton, comprises greywackes, shales, mafic-felsic volcanic rocks along with arenites, limestones, stromatolitic dolomite, carbon phyllites, banded iron formations and banded manganese formations (Harinadha Babu et al., 1981; Chadwick et al., 1988). The rock formations are metamorphosed to greenschist facies and unconformably overlie the older gneiss and mafic/ultramafic enclaves with a basal or near-basal polymictic conglomerate marking the unconformity (Harinadha Babu et al., 1981; Naqvi and Rogers, 1987). The Ranibennur Formation, an upper part of the Chitradurga

Group in the Dharwar-Shimoga greenstone belt is composed of greywacke-argillite, chert and volcanic rocks. The litho units of present study area (Fig. 1) belong to this formation.

The study area comprises a thick sequence of metasediments with greywackes, polymict conglomerate and banded iron formations, with occasional metavolcanic sequences (Fig. 1). In general, the foliation and schistosity of these rocks strike NNW-SSE and their dip varies from 75° to sub-vertical, due east as well as west. The conglomerate, overlying the metavolcanics, occurs as a prominent litho unit in the eastern margin of the belt in the study area. It has pebbles of granite gneiss, quartzite, mafic and felsic volcanic rocks and banded iron formations. Metamorphosed greywackes constitute the bulk of the supracrustal sequences of the northern part of the Dharwar-Shimoga greenstone belt and are best exposed in the quarries at Mandihal and Someshwar near Dharwad. Some outcrops are also located at Uppin Betageri, Gudikatti, Anigol and Bailhongal. The banded iron formations occupy ridges with a general NNW-SSE trend. These iron formations occur interlayered with the thick sequences of greywackes and are in places intensely deformed and brecciated. The metavolcanics occur as flows, pillowed and pyroclastic rocks (volcanic breccia). Based on the geochemical characteristics, Ugarkar et al., (2012, 2014) have classified the flows and pillowed metavolcanics into basaltic komatiites, tholeiitic basalts and tholeiitic basaltic andesites, while the clasts of the pyroclastic breccias are felsic in composition. Doleritic dykes and sills cut across the metavolcanics and metasediments.

PETROGRAPHY

Dharwad greywackes are greenish grey to ash grey in colour, fine to medium grained and in places show the typical shear characteristics of phyllites. They contain sub-angular sand sized mineral grains of quartz, feldspar and deep greenish black rock fragments embedded in a green microcrystalline matrix. Filling of the foliation planes and randomly oriented fracture planes with quartz veinlets (and in some places with calcite) of more than one generation is common.

Petrographically, they are distinctly immature with poorly-sorted angular to sub-angular grains, and generally lack uniformity in the distribution of individual coarse grains. The coarse

grains are largely quartz, plagioclase feldspar and lithic fragments of volcanic (mafic+felsic) rocks, chert and quartzite, with subordinate biotite, K-feldspar and muscovite (Figs. 2a-d). In general, the plagioclase feldspars derived from the plutonic rocks are of comparatively larger dimension (0.3 to 0.8 mm) than those derived from the volcanic rocks (≤ 0.2 mm). Occasionally the plutonic feldspars are well rounded (Fig. 2a). Felsic volcanic lithic fragments are more than the mafic volcanic lithic fragments. In places, fine chlorite needles and scales penetrate clasts of quartz along grain boundaries, which probably is the result of compaction and diagenesis. Quartz is mostly monocrystalline, although polycrystalline variety is also noted. Polycrystalline quartz, characterized by serrated grain boundaries with individual grain size of 0.4 to 0.2 mm (Fig. 2a), is probably derived from gneisses, while the polycrystalline quartz with individual grain size less than 0.02 mm probably indicates recrystallized chert. The grain boundaries of the quartz, feldspars and lithic fragments are serrated and diffused, which could be due to a short transport time and subsequent diagenetic reactions between grains and groundmass. The groundmass consists of chlorite, sericite, plagioclase, quartz and calcareous material. Serrated and diffused grain boundaries of framework grains indicate formation of secondary matrix. It is difficult to determine whether the recrystallized interstitial material originated as detrital matrix or from labile framework grains (see Cox and Lowe, 1996). It is recognized that development of secondary matrix causes problems in determination of QFL percentage and also in the interpretation of provenance based on QFL scheme (Cox and Lowe, 1996). However, the Dharwar greywackes have an average modal composition of 31% quartz, 22% feldspars (plagioclase dominates), 8% lithic fragments (cherts + quartzites + volcanics) and 23% matrix. The lithic fragments are comparatively larger than mineral clasts (Fig. 2a-d).

SAMPLING AND ANALYTICAL METHODS

Relatively fresh homogeneous representative samples of greywackes from northern part of the Dharwar-Shimoga greenstone belt were collected from quarries, road cuttings and natural outcrops. Weathered and jointed surfaces with quartz/calcite veins were avoided. The sample

chips were reduced to sand size by crushing and were then powdered to ~ 200 mesh in an agate mortar to avoid contamination. Major element concentrations were estimated by X-ray fluorescence (XRF: Phillips MAGIX PRO Model 2440) on pressed pellets at the National Geophysical Research Institute, Hyderabad. Determination of trace elements and REEs was carried out using inductively coupled plasma mass spectrometry (ICP-MS: Perkin Elmer Elan DRC II) after dissolving the finely powdered samples by open digestion method using HF-HNO₃-HClO₄ in Teflon beakers at the National Geophysical Research Institute, Hyderabad, following the procedure of Balaram et al. (1996). Loss on ignition (LOI) was determined by heating 1 gm of finely powdered sample in platinum crucible at temperature range between 550 and 600°C in a muffle furnace.

GEOCHEMISTRY AND PROVENANCE

The greywackes of the present study area are characterized by almost uniform silica (59.78-67.96 wt%; av. 62.58), alkali (4.62-7.35 wt%; av. 5.41) contents and SiO₂/Al₂O₃ (3.71-5.25) ratios (Table 1), and compositionally are comparable to the andesite and dacite (Fig. 3a). Moderately higher contents of Al₂O₃ (12.92 to 16.70 wt%), FeO (5.04 to 9.29 wt%) and MgO (3.2 to 6.58 wt%) are suggestive of abundant content of matrix rich in chlorite, clay minerals and iron oxides. As compared to Ranibennur greywackes (Hegde and Chavadi, 2009), the Dharwar greywackes have higher K₂O, CaO, Zr, Y, Σ REE, Th/Sc, Zr/Cr, La/Sc and lower Sr, Cr, Ni, Sc, Cr/Th values (Table 1). The chondrite normalized patterns of these greywackes are characterized by moderately fractionated REE patterns ($La_N/Yb_N = 8.5$) with moderate to high LREE enrichment ($La_N/Sm_N = 4.2$), with almost flat HREE patterns ($Gd_N/Yb_N = \text{average } 1.5$) and moderate negative Eu anomalies (Fig. 3b).

The intensity of chemical weathering of a source terrain can be quantitatively evaluated by the Chemical Index of Alteration (CIA) of clastic rocks (Nesbitt and Young, 1984). The CIA values for Dharwad greywackes range between 57 and 73 with an average value of 63, indicating a moderate degree of weathering. Climatic conditions in the source region also play an important

role in weathering and so in the maturity of sediment that is formed. When plotted on a SiO_2 -
($\text{Al}_2\text{O}_3+\text{K}_2\text{O}+\text{Na}_2\text{O}$) diagram (Suttner and Dutta, 1986), the composition of the Dharwar
greywackes indicate a source area with arid climatic conditions (Fig. 4a). Recently, Ojakangas et
al (2014) have reported glaciomarine origin for the diamictites of Chitradurga Group which
would imply that during the initial stages of Chitradurga Group sedimentation a cold arid climate
prevailed. The Dharwar greywackes belong to the upper part of the Chitradurga Group, their
moderate CIA values may indicate derivation of detritus from fresh basement rocks and from
nearby volcanic sources. The initial composition of the source rocks can be obtained from the
weathering trend line in the A-CN-K diagram (Fig. 4b) of Nesbitt and Young (1984). The trend
line of Dharwar greywackes trace back to dacite, andesite and granodiorite fields.

Cr and Ni abundances in clastic sedimentary rocks are useful in deciphering the
contribution from mafic and ultramafic components in the provenance (McLennan and Taylor,
1984). Cr (53 ppm) and Ni (36 ppm) contents of Dharwar greywackes are low, which suggest
that the felsic component was dominant in the provenance. Th/Sc ratios are used to discriminate
felsic and mafic sources in the provenance (McLennan and Taylor, 1980). Th/Sc ratios more than
>1 reflect an input of sediments from felsic igneous sources, while ratios <0.8 are indicative of
mafic sources. The Dharwar greywackes have Th/Sc ratios more than 1 (average 1.13), which
suggest that the detritus was derived from a felsic dominated igneous source. The chondrite
normalized patterns of Dharwar greywackes are characterized by moderately fractionated REE
patterns with moderate to high LREE enrichment, with almost flat HREE patterns and small
negative Eu anomalies, suggesting felsic dominated source rocks in the provenance. The relative
contribution of felsic and mafic end members (volcanic as well as plutonic rocks) in the
provenance can be modeled using rare earth elements. For this purpose, the average REE data of
granites and tonalities that occur on either side of the study area, and REE data of dacites and
metabasalts from the study area (Devaraju et al, 2007; Dey et al, 2012; Ugarkar, 2012; Ugarkar et
al, 2014) are used as the end members in the provenance (Table 2). The mixing calculations

1 suggest that the average REE pattern is closely matching with a provenance having 40% dacite,
2 30% granite, 20% basalt and 10% tonalite (Fig. 4c).
3
4

5 **DISCUSSION AND CONCLUSION**

6
7 Petrographic evidence, such as the presence of fresh angular plagioclase and volcanic lithic
8 fragments suggest that the Dharwar greywackes were deposited in a turbidite regime very close
9 to provenance in an unstable environment in which erosion, transportation and deposition were
10 so rapid that there was very little scope for weathering and sorting of the clastic material. The
11 detritus comprising felsic and mafic volcanics, feldspars and quartz suggest a mixed source in the
12 provenance. Mafic trace elements and immobile element abundances and their ratios are
13 compatible with a felsic dominated provenance. REE mixing calculations indicate 40% dacite,
14 30% granite, 20% basalt and 10% tonalite composition for the provenance.
15
16

17 It is a well-established fact that the sedimentary basins of Western Dharwar Craton
18 received clastic detritus from an evolved continental crust which had witnessed upper
19 granodiorite crustal differentiation (Srinivasan et al, 1989; Srinivasan and Naqvi, 1990;
20 Srinivasan and Naha, 1993; Chadwick et al, 2000). While subduction related tectonic settings
21 like foreland, back-arc foreland, forearc basins have been suggested by various workers for the
22 deposition of greywackes of the Western Dharwar Craton, (Naqvi et al, 1988; Srinivasan et al,
23 1989; Ugarkar and Nyamati, 2002, Hegde and Chavadi, 2009; Devaraju et al, 2010). When
24 plotted in Th-La-Sc tectonic discrimination diagrams of Bhatia and Crook (1986), greywackes of
25 study area indicate continental island arc tectonic setting for their deposition (Fig. 5a). To
26 distinguish between back-arc foreland and forearc basins, the greywackes were plotted in Hf-Th-
27 Co diagram (Fig. 5b; adopted from Naqvi et al, 2002), wherein the plots spread close to forearc
28 basin than a continental margin basin, indicating a possible sedimentation in a subduction related
29 forearc basin.
30
31

32 In most tectonic settings where stable crust or recycled sediments are exposed, Th/Sc
33 values are more than 1.0. The Th/Sc ratio of the studied samples (av. 1.13) suggests a source area
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

with a predominant granodioritic component and a relatively minor contribution from mafic igneous rocks. The Archaean Western Dharwar Craton was highly evolved at the time of Dharwar sedimentation, as indicated by La/Sc ratios in the greywackes of study area ranging from 2.8-5.8 (av. 4.07), well in excess of upper continental crust (2.2, Taylor and McLennan, 1985). La/Sc ratios in greywackes of study area are also high compared to their ratios in the Ranibennur greywackes, located about 100 km south of Dharwad in the greenstone belt (1.79, Hedge and Chavadi, 2009), suggesting that the greywackes of the northern part of the basin received detritus from more evolved continental crust than the greywackes of the central part of the Dharwar-Shimoga basin. Further, the La/Sc ratio for greywackes of Archaean age from the Canadian Shield (1.45; Camiré et al., 1993), Kalgoorlie, Australia (1.06; Taylor and McLennan, 1985), Fig Tree, Barberton, South Africa (0.54; Hofmann, 2005) and for shales from Zimbabwe (0.94; Fedo et al., 1996) are all significantly less than the greywackes of the study area (av. 4.07). These features indicate that the Archaean crust in the northern part of the WDC was markedly more evolved compared to all the above well-known Archaean cratons.

References

- 1 ABSAR, N. and SREENIVAS, B. (2015). Petrology and geochemistry of greywackes of the ~1.6
2
3 Ga Middle Aravalli Supergroup, northwest India: evidence for active margin processes. Intl.
4
5 Geol. Rew., v.57, pp.134-158.
6
7
8 BALARAM, V., RAMESH, S.L. and ANJANIAH, K.V. (1996). New trace element and REE data in
9
10 thirteen GSF reference samples by ICP-MS. Geostand. Geoanalytical Res., v.20, pp.71–78.
11
12
13 BHATIA, M.R. and CROOK, K.A.W. (1986). Trace element characteristics of graywackes and
14
15 tectonic setting discrimination of sedimentary basins. Contrib. to Mineral. Petrol., v.92,
16
17 pp.181–193.
18
19
20 CAMIRÉ, G.E., LAFLÈCHE, M.R. and LUDDEN, J.N. (1993). Archaean metasedimentary rocks
21
22 from the northwestern Pontiac Subprovince of the Canadian shield: chemical
23
24 characterization, weathering and modelling of the source areas. Precambrian Res., v.62,
25
26 pp.285–305.
27
28
29
30 CHADWICK, B., VASUDEV, V.N. and JAYARAM, S. (1988). Stratigraphy and structure of Late
31
32 Archaean Dharwar volcanic and sedimentary rocks and their basement in a part of the
33
34 Shimoga basin, east of Bhadravathi, Karnataka. J. Geol. Soc. India, v.32, pp.1–19.
35
36
37 CHADWICK, B., VASUDEV, V.N. and HEGDE, G.V. (2000). The Dharwar craton, southern India,
38
39 interpreted as the result of Late Archaean oblique convergence. Precambrian Res., v.99,
40
41 pp.91–111.
42
43
44 COX, R. and LOWE, D.R. (1996). Quantification of the effects of secondary matrix on the
45
46 analysis of sandstone composition, and a petrographic-chemical technique for retrieving
47
48 original framework grain modes of altered sandstones. J. Sed. Res., v.66, pp.548-558.
49
50
51 DEVARAJU, T.C., HUHMA, H., SUDHAKARA, T.L., KAUKONEN, R.J. and ALAPIETI, T.T. (2007).
52
53 Petrology, geochemistry, model and Sm-Nd ages and petrogenesis of the granitoids of the
54
55 northern block of the western Dharwar Craton. J. Geol. Soc. India, v.70, pp.889–911.
56
57
58 DEVARAJU, T.C., SUDHAKARA, T.L., KAUKONEN, R.J., VILJOEN, R.P., ALAPIETI, T.T., AHMED,
59
60 S.A. and SIVAKUMAR, S. (2010). Petrology and geochemistry of greywackes from Goa-
61
62
63
64
65

Dharwar sector, western Dharwar Craton: Implications for volcanoclastic origin. J. Geol. Soc. India, v.75, pp.465–487.

FENG, R. and KERRICH, R. (1990). Geochemistry of fine-grained clastic sediments in the Archean Abitibi greenstone belt, Canada: Implications for provenance and tectonic setting. Geochim. Cosmochim. Acta, v.54, pp.1061–1081.

HARINADHA BABU, P., PONNUSWAMY and KRISHNAMURTHY, K. V. (1981). Shimoga belt. In early Precambrian Supracrustals of Southern Karnataka. Geol. Soc. India Memoir 112, pp.199–218.

HEGDE, V.S. and CHAVADI, V.C. (2009). Geochemistry of late Archaean metagreywackes from the Western Dharwar Craton, South India: Implications for provenance and nature of the Late Archaean crust. Gondwana Res., v.15, pp.178–187.

HOFMANN, A. (2005). The geochemistry of sedimentary rocks from the Fig Tree Group, Barberton greenstone belt: Implications for tectonic, hydrothermal and surface processes during mid-Archaean times. Precambrian Res., v.143, pp.23–49.

LE BAS, M.J., LE MAITRE, R.W., STRECKEISEN, A. and ZANETTIN, B. (1986). A chemical classification of volcanic rocks based on the total alkali-silica diagram. Jour. Petrol., v.27, pp.745–750.

MANIKYAMBA, C., NAQVI, S.M., MOEEN, S., GNANESHWAR RAO, T., BALARAM, V., RAMESH, S.L., AND REDDY, G.L.N. (1997). Geochemical heterogeneities of metagraywackes from the Sandur schist belt: Implications for active plate margin processes: Precambrian Res., v. 84, pp. 117–138.

MCLENNAN, S.M. and TAYLOR, S.R. (1980). Th and U in sedimentary rocks: crustal evolution and sedimentary recycling. Nature, v.285, pp.621–624.

MCLENNAN, S.M. and TAYLOR, S.R. (1984). Archaean sedimentary rocks and their relation to the composition of the Archean continental crust. In: Kroner, A., Hanson, G.N., Goodwin, A.M. (Eds.), Archaean Geochemistry. Springer-Verlag.

NAQVI, S.M. and ROGERS, J.J.W. (1987). Precambrian Geology of India. Clarendon Press,

Oxford.

1 NAQVI, S.M., SAWKAR, R.H., SUBBA RAO, D. V, GOVIL, P.K. and GNANESWAR RAO, T.
2
3 (1988). Geology, geochemistry and tectonic setting of Archaean greywackes from
4
5 Karnataka nucleus, India. *Precambrian Res.*, v.39, pp.193–216.
6

7
8 NAQVI, S.M., UDAY RAJ, B., SUBBARAO, D.V., MANIKYAMBA, C., NIRMAL CHARAN,
9
10 S., BALARAM, V. and SARMA D.S. (2002) Geology and geochemitry of arenite-
11
12 quartzwacke from the Late Archaean Sandur schist belt: Implications for provenance and
13
14 accretion processes. *Precambrian Res.*, v.114, pp.177-197.
15

16
17 NESBITT, H.W. and YOUNG, G.M. (1984). Prediction of some weathering trends of plutonic and
18
19 volcanic rocks based on thermodynamic and kinetic considerations. *Geochim. Cosmochim.*
20
21 *Acta*, v.48, pp.1523–1534.
22

23
24 OJAKANGAS, R.W., SRINIVASAN, R., HEGDE, V.S., CHANDRAKANT, S.M. and SRIKANTIA, S.V.
25
26 (2014). The Talaya conglomerate: an Archaean (~2.7 Ga) Glaciomarine Formation, Western
27
28 Dharwar Craton, Southern India. *Curr. Sci.*, v.106, pp.387-396.
29

30
31 OOTES, L., DAVIS, W.J., BLEEKER, W. and JACKSON, V.A. (2009). Two distinct ages of
32
33 Neoarchaean turbidites in the Western slave Craton: further evidence and implications for a
34
35 possible back-arc model. *J. Geol.*, v.117, pp.15–36.
36

37
38 SRINIVASAN , R. and NAQVI, S.M. (1990). Some distinctive trends in the evolution of the Early
39
40 Precambrian (Archaean) Dharwar craton, South India. *Precambrian Continental Crust and*
41
42 *its economic resources-Developments in Precambrian Geology*. Elsevier, pp. 245–266.
43

44
45 SRINIVASAN, R. and NAHA, K. (1993). Archean sedimentation in the Dharwar craton, southern
46
47 India. *Proc. Natl. Acad. Sci. India*, v.63, pp.1–13.
48

49
50 SRINIVASAN, R., NAQVI, S.M., UDAY RAJ, B., SUBBA RAO, D.V., BALARAM, V. and
51
52 GNANESHWAR RAO, T. (1989). Geochemistry of the Archaean greywackes from the
53
54 northwestern part of the Chitradurga schist belt, Dharwar Craton, South India - evidence for
55
56 granitoid upper crust in the Archaean. *J. Geol. Soc. India*, v.34, pp-505-516.
57

58
59 SUTTNER, L.J. and DUTTA, P.K. (1986). Alluvial Sandstone Composition and Paleoclimate, I.
60
61
62
63
64
65

Framework Mineralogy. SEPM J. Sediment. Res., v.56(3), pp.329-345.


1 TAYLOR, S.R. and MCLENNAN, S.M. (1985). The Continental Crust: Its Composition and
2
3 Evolution, an examination of the geochemical record preserved in sedimentary rocks.
4
5 Blackwell, London.
6

7
8 UGARKAR, A.G. (2012) Petrography and geochemistry of metavolcanics and metasediments of
9
10 the Northern Part of Dharwar-Shimoga Schist Belt, Western Dharwar Craton. Project
11
12 Completion Report submitted to DST, New Delhi.
13
14

15 UGARKAR, A.G. and NYAMATI, R.C. (2002). Geochemical Characteristics of Archean Clastic
16
17 Metasediments of Gadag Gold Field, Southern India: Implications for Provenance and
18
19 Tectonic Setting. Gondwana Res., v.5, pp.245–255.
20
21

22 UGARKAR, A.G., CHANDAN KUMAR, B. and MANUVACHARI, T.B. (2012). Lithology and
23
24 Geochemistry of Metavolcanics and Metasediments of Northern Part of Dharwar-Shimoga
25
26 Schist Belt, Western Dharwar Craton. Indian Mineral., v.46, pp.162–178.
27
28

29 UGARKAR, A.G., CHANDAN KUMAR, B. and MANUVACHARI, T.B. (2014). Geology and
30
31 Geochemistry of Archaean Metavolcanic Rocks of the Northern Part of the Dharwar-
32
33 Shimoga Greenstone Belt, Dharwar Craton, India: Implications for their Petrogenesis and
34
35 Geodynamic Setting. Indian Mineral., v.48, pp.88–107.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65






Fig. 2




Fig. 3






Fig. 4

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65





Fig. 5

Figure Captions:

- Fig. 1. Detailed geological map of northern part of the Dharwar-Shimoga greenstone belt (modified after Ugarkar et al., 2012)
- Fig. 2. Photomicrographs: a) Plagioclase feldspar (pv-volcanic and pp-plutonic source), polycrystalline (pq) and monocrystalline (mq) quartz grains and flakes of mica (m). b) Lithic fragment of mafic volcanic (at the centre. Also note the bimodality and angularity of the grains, as well as the larger dimensions of the rock fragments compared to quartzo-feldspathic grains. c) Lithic fragment of felsic volcanics. d) Quartzite lithic fragment. pv: volcanic and pp: plutonic source. pq: polycrystalline and mq: monocrystalline quartz grains. m: flakes of mica. fv: felsic and mv: mafic volcanic lithic fragments. qz: quartzite lithic fragment.
- Fig. 3. a) $\text{Na}_2\text{O}+\text{K}_2\text{O}$ vs. SiO_2 diagram of Le Bas et al. (1986). b) Chondrite normalized REE diagrams for Dharwar greywackes (chondrite values from Taylor and McLennan, 1985).
- Fig. 4. a) SiO_2 vs. $\text{Al}_2\text{O}_3+\text{Na}_2\text{O}+\text{K}_2\text{O}$ diagram of Suttner and Dutta (1986). b) Ternary plot of A-CN-K (after Nesbitt and Young, 1984). Solid squares are plots of average compositions of typical rock types, Open circles indicate Dharwar greywackes. c) Chondrite normalized REE pattern of average greywackes of study area and estimated REE mixed provenance after mixing the end members in the proportion of 40% dacite, 30% granite, 20% basalt and 10% tonalite.
- Fig. 5. A) Th-La-Sc and b) Hf-Th-Co tectonic discrimination diagrams of Bhatia and Crook (1986) and Naqvi et al. (2002) respectively.

Table 1. Major, trace and rare earth element analyses of greywackes around Dharwad

	Sample	D18	D18/A	D21	D38	D40	S1	S2	S3	N1	N2	N3	N4	Av	RG*
1	SiO ₂	63.93	60.35	59.78	67.96	67.84	61.97	63.79	62.29	60.08	59.95	59.96	63.04	62.58	63.07
2	TiO ₂	0.54	0.64	0.61	0.52	0.51	0.55	0.51	0.61	0.65	0.71	0.65	0.61	0.59	0.45
3	Al ₂ O ₃	13.39	13.93	14.13	13.29	12.92	16.7	15.62	15.89	13.62	14	14.61	14.45	14.38	13.90
4	MnO	0.17	0.17	0.13	0.06	0.07	0.09	0.12	0.1	0.15	0.15	0.15	0.1	0.12	0.09
5	Fe ₂ O ₃	7.71	8.7	7.5	5.04	5.62	8.92	8.18	8.22	8.1	9.285	8.68	6.68	7.72	8.98
6	CaO	3.59	3.51	5.84	1.77	2.26	1.33	2.59	3.23	4.03	3.69	3	3.9	3.23	1.95
7	MgO	4.3	6.58	5.86	3.2	3.48	4.82	3.42	3.78	4.7	5.37	4.52	4.5	4.54	3.92
8	Na ₂ O	3.91	3.18	2.03	4.95	4.44	2.43	2.67	2.77	3.13	3.32	3.65	3.13	3.3	3.65
9	K ₂ O	1.79	1.85	2.59	2.4	2.19	2.52	2.36	2.34	1.7	1.67	1.38	2.56	2.11	1.50
10	P ₂ O ₅	0.17	0.13	0.14	0.2	0.16	0.16	0.13	0.18	0.15	0.16	0.15	0.15	0.16	0.10
11	Total	99.5	99.04	98.61	99.39	99.49	99.49	99.39	99.41	96.31	98.31	96.75	99.12	98.73	99.99
12	Ba	443	477	41	712	582	467	398	460	382	391	412	588	446	
13	Rb	55	67	94	51	73	80	78	87	56	51	68	70	69	65
14	Sr	283	313	339	262	185	141	146	81	356	352	347	290	258	401
15	Ga	19	14	13	19	20	19	17	21	16	14	17	14	17	
16	Ta	0.65	0.82	0.43	0.57	0.62	0.84	0.59	0.73	0.91	0.91	0.91	0.72	0.73	
17	Nb	8.20	10.47	6.75	7.57	7.64	10.48	7.08	9.33	8.23	7.73	8.23	4.87	8	
18	Zr	503	31	33	608	582	639	554	695	97	91	87	95	335	176
19	Y	27	34	29	25	27	28	21	29	26	28	27	22	27	17.4
20	Cr	15	92	83	9	11	15	14	16	104	110	91	73	53	176
21	Ni	39	31	26	22	28	40	40	45	33	58	29	38	36	121
22	Co	23	26	23	9	16	21	18	20	26	26	25	22	21	24
23	V	95	96	97	81	90	104	100	115	110	99	106	86	98	131
24	Cu	27	39	28	23	23	28	26	23	53	32	35	34	31	62
25	Pb	13	8	5	13	16	15	12	12	10	7	2	12	11	
26	Zn	23	27	34	19	23	27	23	24	53	60	53	51	35	49
27	Cs	1.37	1.33	1.59	1.07	1.03	2.51	2.13	1.29	1.67	1.17	1.67	1.30	1.51	
28	Hf	14	1	1	17	16	18	16	20	2	2	2	2	9	4.15
29	Th	9	9	7	9	9	9	8	11	9	8	8	8	9	7.35
30	U	2.65	1.48	1.14	2.52	3.51	2.93	2.64	3.07	2.24	2.24	1.74	1.57	2.31	2.32
31	Sc	6	13	12	5	5	7	7	7	--	--	--	--	8	13.50
32	La	29.83	36.59	37.27	32.27	31.39	30.69	27.65	33.91	33.34	32.34	35.2	36.93	33.12	23.70
33	Ce	56.62	70.74	67.24	60.21	59.56	59.97	53.26	65.62	63.24	63.57	65.32	68.52	62.82	42.70
34	Pr	6.45	8.29	7.67	7.09	6.96	6.98	6.13	7.60	7.30	7.07	7.56	7.39	7.21	6.40
35	Nd	23.61	30.40	28.45	25.82	24.82	25.50	22.51	27.35	27.11	27.15	28.13	28.57	26.62	21.0
36	Sm	4.29	5.68	5.37	4.63	4.52	4.82	4.17	5.03	5.03	5.06	5.26	5.13	4.92	4.60
37	Eu	1.23	1.64	1.41	1.45	1.35	1.26	1.10	1.32	1.41	1.45	1.54	1.39	1.38	1.00
38	Gd	4.41	5.58	5.09	4.58	4.67	4.88	4.06	5.14	4.67	5.16	4.93	4.98	4.85	4.00
39	Tb	0.78	1.03	0.98	0.76	0.82	0.86	0.68	0.88	0.83	0.81	0.86	0.75	0.84	0.50
40	Dy	4.46	5.96	5.23	4.29	4.71	4.76	3.89	4.98	4.64	4.73	4.86	4.03	4.71	2.90
41	Ho	0.87	1.19	1.02	0.84	0.95	0.95	0.75	1.01	0.92	0.96	0.97	0.79	0.94	
42	Er	2.58	3.08	2.72	2.39	2.64	2.81	2.19	2.94	2.44	2.58	2.57	2.16	2.59	
43	Tm	0.43	0.49	0.44	0.42	0.46	0.51	0.42	0.54	0.38	0.38	0.40	0.32	0.43	0.21
44	Yb	2.59	2.66	2.67	2.53	2.74	3.24	2.76	3.37	2.19	2.29	2.32	2.04	2.62	1.80
45	Lu	0.42	0.41	0.40	0.42	0.45	0.53	0.47	0.57	0.32	0.34	0.34	0.31	0.42	

Sample locations: D-around Dharwad, N-north of Dharwad and S-south of Dharwad. *RG – Average values of Ranibennur Greywackes (after Hegde and Chavadi, 2009)

Table 2. REE mixing calculations taking 40% dacite (D), 30% granite (G), 20% basalt (B) and 10% tonalite (T).

	D	B	T	G	Mix*	Gw**
La	24.53	7.54	24.20	32.8	23.59	33.12
Ce	59.98	18.78	26.30	72.7	52.18	62.82
Pr	7.14	2.34	2.61	7.8	5.92	7.21
Nd	30.79	12.50	9.65	22.4	22.50	26.62
Sm	7.87	3.19	2.60	4.2	5.31	4.92
Eu	1.80	0.99	0.39	0.9	1.24	1.38
Gd	7.16	3.98	1.86	2.9	4.72	4.85
Tb	1.45	0.67	0.23	0.5	0.87	0.84
Dy	7.97	3.81	0.75	2.9	4.88	4.71
Ho	1.52	0.75	0.12	0.4	0.90	0.94
Er	4.64	2.27	0.34	1.2	2.70	2.59
Yb	4.75	2.08	0.25	1.1	2.68	2.62
Lu	0.68	0.44	--	0.2	0.41	0.42

Provenance sources: Dacite (Ugarkar, 2012), granite (Devaraju et al, 2007), basalt (Ugarkar et al, 2014) and tonalite (Dey et al, 2012).